



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/66, 31/675		A1	(11) International Publication Number: WO 97/02037
			(43) International Publication Date: 23 January 1997 (23.01.97)
(21) International Application Number: PCT/EP96/02842		S.A., 243, route des Fayards, CH-1290 Versoix (CH). GUYON-GELLIN, Yves [FR/CH]; Symphar S.A., 243, route des Fayards, CH-1290 Versoix (CH). IFE, Robert, John [GB/GB]; SmithKline Beecham Pharmaceuticals, The Frythe, Welwyn, Hertfordshire AL6 9AR (GB). (74) Agent: CONNELL, Anthony, Christopher, SmithKline Beecham, Corporate Intellectual Property, SB House, Great West Road, Brentford, Middlesex TW8 9BD (GB). (81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 26 June 1996 (26.06.96)			
(30) Priority Data: 1920/95-1 30 June 1995 (30.06.95) CH			
(71) Applicants (for all designated States except US): SYMPHAR S.A. [CH/CH]; 243, route des Fayards, CH-1290 Versoix (CH). SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).			
(72) Inventors; and (75) Inventors/Applicants (for US only): NGUYEN, Lan, Mong [CH/CH]; Symphar S.A., 243, route des Fayards, CH-1290 Versoix (CH). NIESOR, Eric [FR/CH]; Symphar S.A., 243, route des Fayards, CH-1290 Versoix (CH). BENTZEN, Craig, Leigh [US/CH]; Symphar S.A., 243, route des Fayards, CH-1290 Versoix (CH). PHAN, Hieu, Trung [CH/CH]; Symphar S.A., 243, route des Fayards, CH-1290 Versoix (CH). DIEP, Vinh, Van [CH/CH]; Symphar S.A., 243, route des Fayards, CH-1290 Versoix (CH). FLORET, Simon [CH/CH]; Symphar S.A., 243, route des Fayards, CH-1290 Versoix (CH). AZOULAY, Raymond [MA/CH]; Symphar S.A., 243, route des Fayards, CH-1290 Versoix (CH). BULLA, Alexandre [RO/CH]; Symphar			


Published

With international search report.

(54) Title: COMPOUNDS AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM

(57) Abstract

Aminophosphonates alpha substituted by phenol groups of formula (I) have lipoprotein(a) lowering activity.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Compounds and Pharmaceutical Compositions containing them

This invention relates to a new therapeutic use of aminophosphonate compounds for lowering plasma and tissue levels of lipoprotein(a). In particular, this invention provides a new use of aminophosphonate derivatives, for the preparation of pharmaceutical compositions useful in the treatment of diseases or disorders associated with high plasma and tissue concentrations of lipoprotein(a); such as, for instance atherosclerosis, thrombosis, restenosis after angioplasty and stroke. This invention also provides a method for increasing thrombolysis and preventing thrombosis and a method of treatment of restenosis after angioplasty by administering to a patient in need thereof an aminophosphonate compound at a dose effective for lowering plasma and tissue lipoprotein(a) levels. In addition, this invention also provides a group of new aminophosphonate compounds for use in the above mentioned uses and compositions.

Recent epidemiologic studies have shown a strong association between elevated lipoprotein(a) [Lp(a)] plasma levels and the occurrence of coronary heart disease, stroke and peripheral artery disease. Lp(a) is now recognized as an independent risk factor for cardiovascular diseases; in addition its role in promoting thrombosis by decreasing thrombolysis is increasingly acknowledged, see for instance "Lipoprotein(a) as A Risk Factor for Preclinical Atherosclerosis" P.J. Schreiner, J.D. Morrisett, A.R. Sharrett, W. Patsch, H.A. Tyroler, K.Wu and G. Heiss; *Arteriosclerosis and Thrombosis* 13, p. 826-833 (1993); "Detection and Quantification of Lipoprotein(a) in the Arterial Wall of 107 Coronary Bypass Patients" M. Rath, A. Niendorf, T. Reblin, M. Dietel, H.J. Krebber and U. Beisiegel; *Arteriosclerosis* 9, p. 579-592 (1989); and "Lipoprotein(a) : Structure, Properties and Possible Involvement in Thrombogenesis and Atherogenesis" A.D. MBewu and P.N. Durrington; *Atherosclerosis* 85, p. 1-14 (1990).

The potential of thrombosis involvement in vessel occlusion and acute cardiovascular syndrome is being increasingly recognized. One of the mechanisms that mediate thrombosis associated with atherosclerotic plaque rupture involves elevated levels of lipoprotein(a). The structure of Lp(a) consists of a low-density lipoprotein (LDL)-like particle with a glycoprotein, apolipoprotein(a) [apo(a)] that is linked via a disulfide bridge to the apo B-100 moiety of the LDL. Structurally there is striking analogy between apo(a) and plasminogen, the precursor of plasmin which cleaves fibrin to dissolve blood clots. However, unlike plasminogen apo(a) is not a substrate for plasminogen activators. This structural resemblance has led researchers to

postulate and later demonstrate that apo(a) interferes with the normal physiological function of plasminogen, leading to a potential thrombogenic activity of Lp(a) see for instance:

- "Activation of Transforming Growth Factor- β is Inversely Correlated with Three
 5 Major Risk Factors for Coronary Artery Disease : Lipoprotein(a), LDL-Cholesterol and Plasminogen Activator Inhibitor-1", A. Chauhan, N.R. Williams, J.C. Metcalfe, A.A. Grace, A.C. Liu, R.M. Lawn, P.R. Kemp, P.M. Schofield and D.J. Grainger; *Circulation*, Vol 90, No. 4, Part 2, p. I-623 (1994); and
 "Influence of Human Apo(a) Expression on Fibrinolysis *in vivo* in Transgenic
 10 Mice" T.M. Palabrica, A.C. Liu, M.J. Aronovitz, B. Furie, B.C. Furie and R. Lawn; *Circulation*, Vol 90, No. 4, Part 2, p. I-623 (1994).

- On the basis of its suspected thrombogenic activity, Lp(a) has also been implicated in peripheral artery disease, in particular stroke. Recently clinicians have shown that
 15 serum Lp(a) levels were significantly higher in stroke patients than in a reference normal population :

- "Lp(a) Lipoprotein in Patients with Acute Stroke" K. Asplund, T. Olsson, M. Viitanen and G. Dahlen; *Cerebrovasc. Diseases* 1, p. 90-96 (1991).
- 20 Restenosis following percutaneous transluminal angioplasty is a common complication occurring in up to 40% of cases within 3-6 months of the intervention. The main cause for restenosis is believed to be abnormal vascular smooth muscle cell activation and proliferation. The proof that high plasma Lp(a) levels are associated with smooth muscle cell proliferation and activation was established *in vitro* and *in*
 25 *vivo* by the two following studies :
- "Proliferation of Human Smooth Muscle Cells Promoted by Lipoprotein(a)" D.J. Grainger, H. L. Kirschenlohr, J.C. Metcalfe, P.L. Weissberg, D.P. Wade and R.M. Lawn; *Science*, Vol 260, p.1655-1658 (1993);and
 "Activation of Transforming Growth Factor- β is Inhibited by Apolipoprotein (a)
 30 *in vivo*", D.J. Grainger, P.R. Kemp, A.C. Liu, R.M. Lawn and J.C. Metcalfe; *Circulation*, Vol 90, No. 4, Part 2, p. I-623 (1994).


- This observation has led to a hypothesis that associates elevated plasma Lp(a) levels with an increased incidence of restenosis. The hypothesis was confirmed
 35 by the results of a recent clinical study showing that, in patients with high plasma Lp(a) levels, a reduction of Lp(a) levels by more than 50% by LDL-apheresis significantly reduced the restenosis rate; see for instance:

- "Effectiveness of LDL-Apheresis in Preventing Restenosis After Percutaneous Transluminal Coronary Angioplasty (PTCA) : LDL-Apheresis Angioplasty Restenosis Trial (L-ART)" H. Yamaguchi, Y. J. Lee, H. Daida, H. Yokoi, H. Miyano, T. Kanoh, S. Ishiwata, K. Kato, H. Nishikawa, F. Takatsu, Y. Kutsumi, H. Mokuno, N. Yamada and A. Noma; Chemistry and Physics of Lipids, Vol 67/68, p. 399-403(1994).

The above discussion has established the rationale for decreasing plasma Lp(a) in patients at risk with elevated levels (>20-30mg/dl). The Lp(a) concentration in individuals appears to be highly determined by inheritance and is hardly influenced by dietary regimes. Various hormones (i.e. steroid hormones, growth hormones, thyroid hormones) have been shown to regulate plasma levels of Lp(a) in man. Of particular interest, drugs which effectively lower LDL such as the bile acid sequestrant cholestyramine or the HMGCoA reductase inhibitors lovastatin or pravastatin do not affect Lp(a) levels. The drugs of the fibrate family : clofibrate or bezafibrate and the antioxidant drug probucol are equally ineffective. The only drug reported to lower Lp(a) is nicotinic acid. However at the high doses necessary for efficacy (4g/day) nicotinic acid has several serious side-effects which preclude its wide use : flushing, vasodilation and hepatotoxicity. Therefore the medical need to lower elevated Lp(a) plasma levels, an independent risk factor for cardiovascular disease, is still unmet.


In contrast to LDL, Lp(a) exists only in mammals high in the evolutionary scale (humans and non human primates) and is exclusively synthesized by the liver cells. Cynomolgus monkeys possess Lp(a) that is similar to human Lp(a), including possession of the unique apolipoprotein apo(a). This primate offers an experimental opportunity for studying the synthesis of Lp(a) and the role of Lp(a) in atherosclerosis and thrombosis. Primary cultures of cynomolgus monkey hepatocytes have been selected as the *in vitro* test for screening aminophosphonate derivatives of formula (I) for their ability to modulate Lp(a) levels. Prior to screening, this assay system had been validated by testing as reference products nicotinic acid and steroid hormones which are known to lower Lp(a) in man.

The present invention relates to the unexpected discovery that aminophosphonate derivatives are effective for lowering plasma and tissue lipoprotein(a). Accordingly, in a first aspect, the present invention provides for the use of a compound of formula (I) :



where:

- X^1, X^2 , which may be identical or different, are H, a straight or branched alkyl or alkoxy group having from 1 to 8 carbon atoms, a hydroxy group or a nitro group,
- 5 X^3 is H, an alkyl group from 1 to 4 carbon atoms, X^3O and one of the two other substituents X^1 or X^2 may form an alkylidene dioxy ring having from 1 to 4 carbon atoms,
- R^1, R^2 , identical or different, are H, a straight or branched alkyl group having from 1 to 6 carbon atoms,
- 10 B is CH_2, CH_2-CH_2 or $CH=CH$,
 n is zero or 1,
- Z is H, a straight or branched alkyl group having from 1 to 8 carbon atoms, an acyl group R^3-CO where R^3 is an alkyl group from 1 to 4 carbon atoms, a perfluoroalkyl group from 1 to 4 carbon atoms,
- 15 A is H, $CH_2-CH=CH_2$, a straight, branched or cyclic alkyl group having from 1 to 8 carbon atoms, or is selected from the following groups :




- where k is an integer from 2 to 4, m is 0 or an integer from 1 to 5, X^4 , X^5 , X^6 , identical or different, are H, a straight or branched alkyl or alkoxy group from 1 to 8 carbon atoms, a hydroxy, trifluoromethyl, nitro, amino, dimethylamino, diethylamino group, a halogen atom (F, Cl, Br, I), X^4 and X^5 may form an alkylidendioxy ring having from 1 to 4 carbon atoms, X^7 is H or CH_3 , R is a straight

or branched alkyl group having from 1 to 6 carbon atoms, an aryl or arylalkyl group from 6 to 9 carbon atoms;
 or a pharmaceutically acceptable salt thereof;
 in the manufacture of a medicament for lowering plasma and tissue lipoprotein(a).

5

European Patent Application EP 0'559'079A (1993) [corresponding to the US Patent 5 424 303] discloses compounds of formula (I) as well as their use in decreasing plasma cholesterol and blood peroxides.

- 10 Preferred compounds of formula (I) for use in the manufacture of a medicament for lowering plasma and tissue lipoprotein(a) are those of the formula (Ia):



- where B, R¹, R², X¹, X², X³, X⁴, Z, n and m are as hereinbefore defined;
 15 or a pharmaceutically acceptable salt thereof.

Certain compounds within the scope of formula (Ia) are novel and are particularly useful in lowering plasma and tissue lipoprotein(a).

- 20 Accordingly, in a further aspect, this invention provides aminophosphonate derivatives of formula (Ia) where:
 X¹ is H, C₍₁₋₈₎alkyl or C₍₁₋₈₎alkoxy;
 X² is C₍₁₋₈₎alkyl or C₍₁₋₈₎alkoxy;
 X³ is H, C₍₁₋₄₎alkyl, or X³O and one of the two other substituents X¹ or X² may
 25 form an alkylidene dioxy ring having from 1 to 4 carbon atoms;
 R¹, R², which may be identical or different, are H or C₍₁₋₆₎alkyl;
 B is CH₂-CH₂, CH=CH or CH₂;
 n is zero or 1;
 Z is H or C₍₁₋₈₎alkyl;
 30 m is an integer from 0 to 5;
 X⁴ is H, C₍₁₋₈₎alkyl, C₍₁₋₈₎alkoxy, or halo;

- and the pyridyl ring is attached by the ring carbon α - or β - to the nitrogen (2- or 3-pyridyl);
- or a salt, preferably a pharmaceutically acceptable salt, thereof; and excluding:
- Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-pyridyl)
- 5 aminomethylphosphonate;
- Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(2-picolyl)
- aminomethylphosphonate;
- Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-picolyl)
- aminomethylphosphonate;
- 10 Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-methyl-N-(3-picolyl)
- aminomethylphosphonate;
- Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(2-pyridylethyl)
- aminomethylphosphonate, and
- Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(4-picolyl)
- 15 aminomethylphosphonate.
- Suitably, X^1 is H, $C_{(1-4)}$ alkyl or $C_{(1-4)}$ alkoxy, preferably $C_{(1-3)}$ alkyl or $C_{(1-3)}$ alkoxy, more preferably hydrogen, methyl or methoxy.
- 20 Suitably, X^2 is $C_{(1-4)}$ alkyl or $C_{(1-4)}$ alkoxy, preferably $C_{(1-3)}$ alkyl or $C_{(1-3)}$ alkoxy, more preferably methyl or methoxy.
- Suitably, X^1 and X^2 are both alkoxy or one of X^1 and X^2 is alkyl and the other is alkoxy, or one of X^1 and X^2 is $C_{(1-4)}$ alkyl and the other of X^1 and X^2 is $C_{(1-3)}$ alkyl.
- 25 Suitably combinations of X^1 and X^2 include methoxy and methoxy, methoxy and methyl, n-propyl or iso-butyl, methyl and methyl or t-butyl, respectively.
- 30 Preferably, X^3 is hydrogen.
- Preferably, $(B)_n$ is a direct bond.
- Preferably, R^1 and R^2 is each a $C_{(1-3)}$ alkyl group, more preferably, a C_2 or C_3 alkyl group, in particular R^1 and R^2 is ethyl or isopropyl.
- 35 Preferably, Z is hydrogen.
- Preferably, X^4 is hydrogen or methyl which is preferably on the ring carbon adjacent to N.
- 40 Preferably, the pyridyl ring is attached by the ring carbon β - to the nitrogen (3-pyridyl).

When used herein, the terms 'alkyl' and 'alkoxy' include both straight and branched groups, for instance, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, s-butyl, t-butyl, etc..

5

Preferred compounds of formula (Ia) include:

Diisopropyl α -(4-hydroxy-3-methoxy-5-methylphenyl)-N-(3-pyridyl)-aminomethylphosphonate;

10 Diisopropyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate;

Diethyl α -(3-methyl-4-hydroxy-5-t-butylphenyl)-N-(3-pyridyl)-aminomethylphosphonate;

Diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate; and

15 Diethyl α -(3,5-dimethyl-4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate.

Independently from the previously published activity, the present invention relates to the unexpected discovery that aminophosphonate derivatives of formula (I) are effective for decreasing Lp(a) production by primary cultures of Cynomolgus monkey hepatocytes. Lp(a) of these primates is similar in immunologic properties to human Lp(a) and occurs in an almost identical frequency distribution of plasma concentrations, see for instance:

20 "Plasma Lipoprotein(a) Concentration is Controlled by Apolipoprotein(a) Protein Size and the Abundance of Hepatic Apo(a) mRNA in a Cynomolgus Monkey Model", N. Azrolan, D. Gavish and J. Breslow; J. Biol. Chem., Vol 266, p. 13866-13872 (1991).

Therefore the compounds of this invention are potentially useful for decreasing Lp(a) in man and thus provide a therapeutic benefit.

30

In particular, this invention provides a new therapeutic use for aminophosphonate compounds of formula (I) as Lp(a) lowering agents. Diseases associated with elevated plasma and tissue levels of lipoprotein(a) include, for instance, coronary heart disease, peripheral artery disease, intermittent claudication, thrombosis, restenosis after angioplasty, extracranial carotid atherosclerosis, stroke and atherosclerosis occurring after heart transplant.

35

The recently discovered Lp(a) lowering activity of the aminophosphonates of formula (I) is independent from their previously reported pharmacological activities of decreasing plasma cholesterol and blood peroxides. Recent clinical studies have shown that neither the hypocholesterolemic drug pravastatin nor the antioxidant drug probucol can decrease Lp(a) levels in man. See for example :

- 5 "Serum Lp(a) Concentrations are Unaffected by Treatment with the HMG-CoA Reductase Inhibitor Pravastatin: Results of a 2-Year Investigation" H.G. Fieseler, V.W. Armstrong, E. Wieland, J. Thiery, E. Schütz, A.K. Walli and D. Seidel; Clinica Chimica Acta, Vol 204, p. 291-300 (1991); and
- 10 "Lack of Effect of Probucol on Serum Lipoprotein(a) Levels", A. Noma; Atherosclerosis 72, p. 267-269 (1989).

- For therapeutic use the compounds of the present invention will generally be administered in a standard pharmaceutical composition obtained by admixture with a pharmaceutical carrier selected with regard to the intended route of administration and standard pharmaceutical practice. For example, they may be administered orally in the form of tablets containing such excipients as starch or lactose, or in capsule, ovules or lozenges either alone or in admixture with excipients, or in the form of elixirs or suspensions containing flavouring or colouring agents. They may be
- 15 injected parenterally, for example, intravenously, intramuscularly or subcutaneously. For parenteral administration, they are best used in the form of a sterile aqueous solution which may contain other substances, for example, enough salts or glucose to make the solution isotonic with blood. The choice of form for administration as well as effective dosages will vary depending, inter alia, on the condition being treated.
- 20 The choice of mode administration and dosage is within the skill of the art.
- 25

The compounds of structure (I) and their pharmaceutically acceptable salts which are active when given orally can be formulated as liquids, for example syrups, suspensions or emulsions or as solids for example, tablets, capsules and lozenges.

- 30 A liquid formulation will generally consist of a suspension or solution of the compound or pharmaceutically acceptable salt in a suitable liquid carrier(s) for example, ethanol, glycerine, non-aqueous solvent, for example polyethylene glycol, oils, or water with a suspending agent, preservative, flavouring or colouring agents.
- 35 A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations. Examples of such carriers include magnesium stearate, starch, lactose, sucrose and cellulose.

A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatin capsule; alternatively, a dispersion or suspension can be prepared using any suitable pharmaceutical carrier(s), for example aqueous gums, celluloses, silicates or oils and the dispersion or suspension then filled into a soft gelatin capsule.

Typical parenteral compositions consist of a solution or suspension of the compound or pharmaceutically acceptable salt in a sterile aqueous carrier or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilised and then reconstituted with a suitable solvent just prior to administration.

A typical suppository formulation comprises a compound of structure (I) or a pharmaceutically acceptable salt thereof which is active when administered in this way, with a binding and/or lubricating agent such as polymeric glycols, gelatins or cocoa butter or other low melting vegetable or synthetic waxes or fats.



Preferably the composition is in unit dose form such as a tablet or capsule.

Each dosage unit for oral administration contains preferably from 1 to 250 mg (and for parenteral administration contains preferably from 0.1 to 25 mg) of a compound of the structure (I) or a pharmaceutically acceptable salt thereof calculated as the free base.

The pharmaceutically acceptable compounds of the invention will normally be administered to a subject in a daily dosage regimen. For an adult patient this may be, for example, an oral dose of between 1 mg and 500 mg, preferably between 1 mg and 250 mg, or an intravenous, subcutaneous, or intramuscular dose of between 0.1 mg and 100 mg, preferably between 0.1 mg and 25 mg, of the compound of the structure (I) or a pharmaceutically acceptable salt thereof calculated as the free base, the compound being administered 1 to 4 times per day.

Compounds of formula (I) may be prepared according to the processes described in European Patent Application EP 0 559 079-A (1993)[corresponding to the US Patent 5


424 303]. This process which has two variants is shown in the following general scheme:

GENERAL SYNTHESIS SCHEMEVariant 1Variant 2

Variant 1 is used when Z is H, i. e. when the starting compound is a primary amine. Briefly, the aminophosphonates of formula (I) are prepared by nucleophilic addition of a dialkyl phosphite or its sodium salt obtained *in situ* by the reaction of dialkyl phosphite and sodium hydride on the imine obtained by condensation of the
 5 appropriate aldehyde and a primary amine.

Variant 2 is used when Z is not H, i. e. when the starting compound is a secondary amine. In this case, the aminophosphonates of formula (I) are prepared by reacting
 10 equimolar amounts of the appropriate aldehyde and the secondary amine and a dialkyl phosphite. The reaction is advantageously carried out in the presence of p-toluenesulfonic acid as a catalyst in a hydrocarbon solvent such as benzene or toluene with concomitant elimination of water, for instance, by using a Dean-Stark
 apparatus.

15 Novel compounds of formula (Ia) in which Z is hydrogen may be prepared by a process which comprises treating an imine of formula (II):



20 (II)

in which B, X¹, X², X³, X⁴, m and n are as hereinbefore defined;
 with a phosphite compound of formula (III):



25 (III)


in which R¹ and R² are as hereinbefore defined; or a trialkyl silyl derivative thereof, preferably the trimethyl silyl phosphite, or a metal salt thereof, for instance the sodium salt, formed *in situ* by treatment of the compound of formula (III) with a
 suitable base, for instance sodium hydride, ethoxide or methoxide

30

The reaction may be carried out in the presence or absence of a catalyst. Suitable catalysts include amine such as diethylamine or triethylamine. The reaction may be carried out in the absence or presence of a solvent. Suitable solvents include

petroleum ether, benzene, toluene, diethyl ether, tetrahydrofuran, 1,2-dimethoxyethane. Suitable reaction temperatures are in the range 30 to 140°C.

The imine compound of formula (II) may be obtained by condensing an aldehyde compound of formula (V):



(V)

in which B, X¹, X², X³ and n are as hereinbefore defined;
with a primary amine of formula (VI):



(VI)

in which A is as as hereinbefore defined;
under imine forming conditions.

Suitably, the condensation may be effected with or without a catalyst in a solvent such as ether, tetrahydrofuran, benzene, toluene or ethanol. Suitable catalysts include molecular sieve, an acid such as glacial acetic acid, p-toluene sulphonic acid, thionyl chloride, titanium tetrachloride, boron trifluoride etherate, or a base such as potassium carbonate.. The reaction is suitably carried out at a temperature in the range 0°C to the boiling point of the solvent being used. For less reactive amines/aldehydes, the reaction may be usefully carried out in a Dean-Stark apparatus.

Novel compounds of formula (Ia) in which Z is not hydrogen may be prepared by a process which comprises treating equimolar amounts of an aldehyde of formula (V), a secondary amine of formula (VII):




(VII)

in which Z is a C₍₁₋₈₎alkyl group and A is as hereinbefore defined; and a phosphite of formula (III), suitably in the presence of p-toluenesulfonic acid as a catalyst, in a hydrocarbon solvent such as petroleum ether, benzene, toluene or xylene, at a temperature between ambient temperature and the boiling point of the


solvent being used, and with concomitant elimination of water, for instance, by using a Dean-Stark apparatus.

- Compounds of formula (Ia) in which m is not zero may also be prepared by a process which comprises treating a compound of formula (VIII):



(VIII)

- 10 in which B, R¹, R², X¹, X², X³ and n are as hereinbefore defined; an aldehyde of formula (IX):



(IX)

- 15 in which m is an integer from 1 to 5 and X⁴ is as hereinbefore defined; under reductive amination conditions.

- Suitable such conditions include carrying out the reaction in the presence of sodium cyanoborohydride in an alcoholic solvent, preferably methanol, at a pH between 3 to 6 and at a temperature between 0°C and 25°C.


- A compound of formula (VIII) may be obtained according to the process hereinbefore described for a compound of formula (Ia) from an aldehyde of formula (V), a secondary amine of formula (VII) in which Z is protecting group which can be removed by hydrogenolysis, for instance an α substituted benzyl or benzyloxycarbonyl and a phosphite of formula (III). This forms an intermediate which is then subjected to hydrogenolysis according to standard conditions, to give a compound of formula (VIII).

- 30 Through their amino function, the aminophosphonate ester (I) can form salts of inorganic acids such as HCl, H₂SO₄ or with organic acids such as oxalic acid, maleic

acid, sulfonic acids, etc.. An example of hydrochloride salt of aminophosphonate (I) is provided (example 5). All these salts are integral part of this invention.

Compounds of structure (I) are racemates as they have at least one chiral center which is the carbon atom in position alpha to the phosphonate group. The compounds (I) therefore exist in the two enantiomeric forms. The racemic mixtures (50% of each enantiomer) and the pure enantiomers are comprised in the scope of this application. In certain cases, it may be desirable to separate the enantiomers.

In a further aspect, the present invention provides a process for the enantiomeric synthesis of a derivative of formula (I) which process comprises treating either of the (+) or (-) enantiomer of the α -substituted aminomethylphosphonate of formula (X):



in which B, R¹, R², X¹, X², X³ and n are as hereinbefore defined; (X)

with an aldehyde of formula (XI):



(XI)

in which R³ is as hereinbefore defined; under reductive amination conditions.

Suitable such conditions include carrying out the reaction in the presence of sodium cyanoborohydride in an alcoholic solvent, preferably methanol, at a pH between 3 to 6 and at a temperature between 0°C and 25°C.


The key α -substituted primary aminomethylphosphonate of formula (X) is obtained by treating an aldehyde of formula (V), as hereinbefore defined, with (+) or (-)- α -methylbenzylamine to form an intermediate imine which is then reacted with a phosphite ester HPO(OR¹)(OR²) to give a mixture of diastereoisomers which may be separated by conventional techniques, for instance fractional crystallisation or chromatography. Hydrogenolysis can then be used to remove the benzyl group from nitrogen, to give the α -substituted primary aminomethyl-phosphonate of formula (X). This approach is illustrated by the preparation of enantiomers of compounds No. 7

and 15 of Table 1. Alternately, the resolution of the aminophosphonate racemates can be effected by preparative chiral chromatography, in particular chiral HPLC. The experimental conditions for chromatographic separation of enantiomers of compound No. 20 are provided. With either separation method, final enantiomeric purity can be ascertained by measuring the specific rotations of the separated isomers.

The structure of compounds of formula (I) were established by their elemental analysis, their infrared (IR), mass (MS) and nuclear magnetic resonance (NMR) spectra. The purity of the compounds was checked by thin layer, gas liquid or high performance liquid chromatographies.


The invention is further described in the following examples which are intended to illustrate the invention without limiting its scope. In the tables, n is normal, i is iso, s is secondary and t is tertiary. In the description of the NMR spectra, respectively s is singlet, d doublet, t triplet and m multiplet. TsOH is p-toluenesulfonic acid monohydrate. The temperatures were recorded in degrees Celsius and the melting points are not corrected. In the measurement of optical activity, an enantiomer which rotates the plane of polarized light to the right is called dextrorotatory and is designated (+) or (D). Conversely, levorotatory defines an enantiomer which rotates the plane of polarized light to the left, designated (-) or (L). Unless otherwise indicated, the physical constants and biological data given for aminophosphonates of formula (I) refer to racemates.

Example 1 - Dimethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-pyridyl)-amino-methylphosphonate



- A mixture of 50g (0.206mol) of 3,5-di-tert-butyl-4-hydroxybenzaldehyde and 20.3 g (2.16 mol) of 3-aminopyridine dissolved in 300 ml toluene and a catalytic amount of p-toluenesulfonic acid (ca. 50 mg) contained in a flask connected to a Dean Stark apparatus was refluxed for 17 h. The solution was evaporated to dryness to give a solid which was purified by recrystallisation from ligroin : mp = 125-130°, IR (KBr) : 1590 cm^{-1} : CH=N.
- Dimethyl phosphite (63.8 g, 0.58 mol) was added to 60 g (0.19 mol) of the previously described imine dissolved in 230 ml THF and the mixture was refluxed for 6 h. The solvent was evaporated and the residue was purified by column chromatography (SiO_2 , 9/1 $\text{CHCl}_3/\text{MeOH}$). Recrystallisation from a mixture of methyl-tert-butyl ether/petroleum ether gave a white solid, mp = 168-170°C.
- IR (KBr) = 3300 cm^{-1} : NH, 1240 : P=O, 1030 : P-O-C
 NMR (CDCl_3) : δ = 8.06, 7.96, 7.4 and 6.9 (4m, 1H each) : aromatic H, 3-pyridyl, 7.2 (d, $J_{\text{P-H}} = 2\text{Hz}$, 2H) : aromatic H, substituted phenyl, 5.24 (s, 1H) : OH, 4.66 (d, $J_{\text{P-H}} = 22\text{Hz}$, 1H) : $\text{CH-PO}_3\text{Me}_2$, 4.75 - 4.68 (m, 1H) : NH, 3.74 and 3.39 = (two d, $J = 11\text{Hz}$) : P-O- CH_3 , 1.42 (s, 18H) : tert-Bu
- MS : m/e = 419 : $\text{M}^+ - 1$, 311 (100%) : $\text{M}^+ - \text{PO}_3\text{Me}_2$

Example 2 - Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(2-pyridyl)-amino-methylphosphonate




- The process described in example 1 was employed using 2-aminopyridine as the amine and diethyl phosphite as the phosphonate reagent. The title compound was purified by column chromatography (95/5 $\text{CHCl}_3/\text{MeOH}$) to yield a solid (61%); mp = 116-118° (AcOEt-ligroin)
 MS (m/e) = 448 : M^+ , 311 : $\text{M}^+ - \text{PO}_3\text{Et}_2$, 78 (100%) : $\text{C}_5\text{H}_4\text{N}$

$\delta =$ 8.09, 7.38 and 6.57 and 6.44 (4m, 1H each): aromatic H, 2-pyridyl, 7.28 (d, J_{P-H}=2Hz, 2H): aromatic H, substituted phenyl, 5.46 (dd, J = 9 and 22 Hz, 1H): CH-PO₃Et₂, 5.3 (m, 1H): N-H, 4.14-3.66 (3m, 4H total): P-O-CH₂-CH₃, 1.42 (s, 18H): tert-Bu, 1.21 and 1.16 (2 t, 3H each): P-O-CH₂-CH₃

5

Example 3 - Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-[5-(2-chloro pyridyl)]-aminomethylphosphonate




The process described in example 1 was employed using 5-amino-2-chloropyridine as the amine and diethyl phosphite as the phosphonate reagent. The title compound was obtained in 50% yield after column chromatography (98/2 CHCl₃/MeOH) and trituration in petroleum ether; mp = 124-126°C

MS (m/e) = 483: M⁺ + 1, 345 (100%), 347 (30%): M⁺ - PO₃Et₂

NMR (CDCl₃): $\delta =$ 7.78, 7.05 and 6.09 (3H): aromatic H, 3-pyridyl, 7.18 (d, J=2Hz, 2H): aromatic H, substituted phenyl, 5.22 (s, 1H): OH, 4.83 (t, J=8Hz): N-H, 4.57 (dd, J=7.5 and 22.5Hz): CH-PO₃-Et₂, 4.1, 3.86 and 3.56 (3m, 4H): P-O-CH₂-CH₃, 1.40 (s, 18H): t-Bu, 1.28 and 1.05 (2t, J=7Hz): P-O-CH₂-CH₃


Example 4 - Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-acetyl-N-(4-picolyl)-aminomethylphosphonate



A mixture of acetic anhydride (1.4g, 14 mmol), diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(4-picolyl)-aminomethylphosphonate (6 g, 13 mmol) and triethyl amine (1.9 ml, 14 mmol) in 20 ml toluene was refluxed for 16 h. The reaction mixture was extracted with brine, dried and evaporated to dryness. The residue was recrystallized in a mixture of dichloromethane and petroleum ether to give 3.7 g (57% yield); mp = 160-162°C.

MS (m/e): 504: M⁺, 461: M⁺ - COCH₃, 367: M⁺ - PO₃Et₂, 325 (100%): M⁺ + 1 - PO₃Et₂ - COCH₃

Example 5 - Hydrochloride salt of diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-pyridyl)aminomethylphosphonate




- 5 Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-pyridyl)aminomethylphosphonate (3 g, 6.7 mmol) was dissolved with slight warming in 60 ml toluene and the resulting solution was saturated with gaseous hydrogen chloride. After 16 h at 0°C the mixture was evaporated to dryness and the residue was recrystallized in EtOH; mp = 193-194°C.

10 Elemental analysis: $C_{24}H_{38}ClN_2O_4P$

% Calc.	C 59.43	H 7.90	Cl 7.31	N 5.78	P 6.39
% Found	C 59.53	H 8.10	Cl 7.02	N 5.72	P 6.21

Example 6 - Diethyl α -(3,4-methylenedioxyphenyl)-N-(3-pyridyl)aminomethylphosphonate

15



The process described in example 8 was followed. The title compound was purified by column chromatography (9/1 $CHCl_3/MeOH$); 60% yield, mp = 98-99°C, $C_{17}H_{21}N_2O_5P$.


20 IR (KBr) = 1240 cm^{-1} : P=O, 1030: P-O-C

MS (m/e) = 365 $M^+ + 1$, 227 (100%): $M^+ - PO_3Et_2$

NMR ($CDCl_3$) δ = 8.1, 7.95, 7.05 and 6.95 (4m, 1H each): aromatic H, 3-pyridyl, 6.90, 6.85, 6.75 (3m, 3H): aromatic H, substituted phenyl, 5.95 (2H): =O-CH₂-O, 4.86 (d x d, 1H, J=8 and 10Hz): N-H, 4.63 (d x d, 1H, J=8 and 24 Hz): CH-PO₃Et₂,

25 4.18-3.70 (3m, 4H total): P-O-CH₂-CH₃, 1.31 and 1.16: (2t, J=7Hz): P-O-CH₂-CH₃

Example 7 - Diethyl α -(4-hydroxyphenyl)-N-(3-pyridyl)aminomethylphosphonate




4-Hydroxybenzaldehyde (6 g, 49 mmol) was reacted at room temperature with 3-aminopyridine (4.5 g, 52 mmol) in 30 ml THF at room temperature to give 9.9 g of a light brown solid. The imine so obtained (5.9 g, 30 mmol) was dissolved in 50 ml

- 5 THF, diethyl phosphite was added in two portions, one at the beginning of the reaction and the other after 6 h at reflux, (total amount: 8.2 g, 60 mmol). The reaction mixture was refluxed overnight. Filtration of the precipitate formed gave 7.5 g (75%) of a tan solid, mp = 210-212°C (EtOH).

MS (m/e) = 337 : $M^+ + 1$, 199 (100%): $M^+ - PO_3Et_2$

- 10 NMR (DMSO- d_6): δ = 9.35 (s, 1H): OH, 8.15, 7.7, 7.1 and 7.0 (4m, 1H each): aromatic H, 3-pyridyl, 6.5 (dxd, 1H): N-H, 7.3 and 6.7 (2m, 2H each) : aromatic H, 4-hydroxyphenyl, 4.93 (dxd, 1H) : $CH-PO_3Et_2$, 4.1-3.6 (3m, 4H total): P-O- CH_2 - CH_3 , 1.15 and 1.02 (2t, 3H each) : P-O- CH_2 - CH_3

- 15 Example 8 - Diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-amino-methylphosphonate



A mixture of 3 g (16.4 mmol) of syringaldehyde and 1.63 g (17.3 mmol) of 3-aminopyridine dissolved in 10 ml toluene and a catalytic amount of p-toluenesulfonic acid (ca. 5 mg) contained in a flask connected to a Dean Stark apparatus was refluxed for 17 h. The solution was evaporated to dryness to give 4.2 g (100%) of the crude imine. Diethyl phosphite (4.8 g, 35 mmol) was added to 4.2 g (17.3 mmol) of the previously described imine dissolved in 10 ml THF and the mixture was refluxed for 7 h. Another amount of diethyl phosphite (4.8 g, 35 mmol) was added and the mixture was refluxed overnight (total reaction time : 17 h). The solvent and the excess of diethyl phosphite were evaporated and the residue was recrystallized from a mixture of ethanol and dichloromethane to give 4.2 g (61%) of a white solid, mp = 181-183°.

IR (KBr) = 1240 cm^{-1} : P=O and 1030 : P-O-C

- 30 MS (m/e) = 397 : $M^+ + 1$, 259 (100%) : $M^+ - PO_3Et_2$


NMR (CDCl₃): δ = 8.08, 7.98, 7.04 and 6.84 (4m, 1H each): aromatic H, 3-pyridyl, 6.69 (d, J = 2Hz, 2H): aromatic H, substituted phenyl, 5.8 (broad, 1H): OH, 4.84 (d x d, 1H, J=7 and 10Hz): N-H, 4.62 (d x d, 1H, J=7 and 23 Hz): CH-PO₃Et₂, 4.18-3.65 (3m, 4H total): P-O-CH₂-CH₃, 3.86 (s, 6H): OCH₃, 1.31 and 1.16: (2t, J=7Hz): P-O-

5 CH₂-CH₃

Elemental analysis: C₁₈H₂₅N₂O₆P

% Calc.	C 54.54	H 6.36	N 7.07	P 7.81
% Found	C 54.50	H 6.38	N 6.99	P 7.65

10 Example 9 - Diethyl α -(3,4,5-trimethoxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate




A mixture of 3,4,5-trimethoxybenzaldehyde (10 g, 51 mmol) and 3-aminopyridine (4.8 g, 51 mmol) and a catalytic amount of TsOH in 50 ml toluene was refluxed for 16 h in a flask connected to a Dean-Stark trap. Evaporation of toluene gave 12.9 g (93%) of the crude imine which was used directly in the next reaction.

A 50 ml THF mixture containing the imine (6g, 22 mmol) and diethyl phosphite (6.1g introduced at the beginning and 6.1 g after 4 h, total amount = 12.2 g, 88 mmol) was refluxed for 8 h. The residue after evaporation of THF and excess of HPO₃Et₂ was triturated in petroleum ether to give 7.12 g (79%) of a white solid, mp = 135-137°C.

20 MS (m/e) = 410 : M⁺, 273 (100%) : M⁺ - PO₃Et₂

NMR (CDCl₃): δ = 8.1, 8.0, 7.05 and 6.85 (4m, 1H each): aromatic H, 3-pyridyl, 6.69 and 6.68: (d, J = 2Hz, 2H): aromatic H, substituted phenyl, 4.86 (d x d, 1H, J=8 and 10Hz): N-H, 4.63 (d x d, 1H, J=7 and 23 Hz): CH-PO₃Et₂, 4.18-3.70 (3m, 4H total): P-O-CH₂-CH₃, 3.86 (two s, 9H): OCH₃, 1.31 and 1.16: (2t, J=7Hz): P-O-CH₂-CH₃

Example 10 - Diethyl α -(3-ethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate




5 A 50 ml toluene solution containing 3-ethoxy-4-hydroxybenzaldehyde (10 g, 60 mmol), 3-aminopyridine (5.6 g, 60 mmol) and 50 mg of TsOH placed in a flask connected to a Dean-Stark trap was refluxed for 4 h to give 14.62 g (95%) of the corresponding imine.

To a suspension of sodium hydride (1.19 g of a 60% mixture, 30 mmol) in 20 ml dry THF was added HPO_3Et_2 (9.12 g, 66 mmol) under nitrogen and the resulting mixture was stirred until the initial turbid suspension became completely clear. To this solution of NaPO_3Et_2 was added the above imine (8 g, 33 mmol) dissolved in 10 ml THF and the resulting solution was refluxed for 2 h. THF was evaporated and the residue was partitioned into H_2O and CH_2Cl_2 . Evaporation of the dried organic phase gave 3.1 g of a white solid, mp = 184-187°C.

MS : (m/e) = 380 : M^+ , 243: $\text{M}^+ - \text{PO}_3\text{Et}_2$

15 NMR (DMSO- d_6): δ = 8.9 (s, 1H): OH, 8.15, 7.3, 7.0 and 6.9 (1H each): aromatic H, 3-pyridyl, 7.1 (m, 2H) and 6.68 (d, J = 8 Hz, 1H): aromatic H, phenyl, 6.5 (dxd, J = 6 and 10 Hz): NH, 4.92 (dxd, J = 10 and 24 Hz): $\text{CH}-\text{PO}_3\text{Et}_2$, 4.05-3.6 (4m, 6H total): P-O- $\text{CH}_2\text{-CH}_3$ and OCH_2CH_3 , 1.29 (t, J = 7Hz, 3H): O- $\text{CH}_2\text{-CH}_3$, 1.16 and 1.04 (2t, J = 7Hz, 3H each): P-O- $\text{CH}_2\text{-CH}_3$

20 Example 11 - Diethyl α -(4-hydroxy-3-methoxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate




The procedure described in example 10 was followed, using 4-hydroxy-3-methoxybenzaldehyde as the starting material. The title compound is a white solid, mp = 170-173°C.

MS (m/e) = 366: M^+ , 229: $\text{M}^+ - \text{PO}_3\text{Et}_2$

NMR (DMSO- d_6) δ = 8.9 (s, 1H): OH, 8.15, 7.75, 7.0 and 6.9 (4m, 4H): aromatic H, 3-pyridyl, 7.1 (m, 2H) and 6.7 (d, J = 8 Hz, 1H): aromatic H, phenyl, 6.5 (dxd, J = 6


and 10 Hz): NH, 4.92 (dxd, J = 10 and 24 Hz): CH-PO₃Et₂, 4.05-3.6 (3m, 4 H total): P-O-CH₂-CH₃, 3.72 (s, 3H): OCH₃, 1.17 and 1.4 (2t, J = 7Hz, 6H): P-O-CH₂-CH₃

Example 12 - Diethyl α-(3,5-dimethoxy-4-hydroxyphenyl)-N-(4-picolyl)-aminomethylphosphonate



- A solution of 2.5 g (13.7 mmol) syringaldehyde and 1.6 g (14.4 mmol) 4-picolylamine dissolved in 100 ml toluene contained in a flask connected to a Dean-Stark apparatus was refluxed for 3 h. Toluene was evaporated under vacuum then the residue dissolved in 10 ml THF was heated with 5.1 g (36.8 mmol) diethyl phosphite for 6 h. THF was evaporated and the residue was purified by column chromatography (SiO₂, 95/5 CHCl₃/MeOH). Recrystallisation in a mixture of CH₂Cl₂-petroleum ether gave 3.7 g (45%) of a solid, mp = 124-126°C.
- MS (m/e) = 410: M⁺, 273: M⁺-PO₃Et₂
- NMR (CDCl₃) δ = 8.55 and 7.22 (2m, 4H): aromatic H, 4-picolyl, 6.75 (d, J = 2Hz, 2H): aromatic H, phenyl, 4.15-3.77 (several m, 5 H): P-O-CH₂-CH₃ and CH-PO₃Et₂, 3.89 (s, 6H): OCH₃, 3.82 and 3.62 (2d, J = 14 Hz): NH-CH₂-Py, 1.33 and 1.16 (2t, J = 7Hz, 6H): P-O-CH₂-CH₃


Example 13 - Diethyl α-(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-picolyl)-aminomethylphosphonate



- The procedure described in example 12 was followed, using 3-picolylamine as the starting material. The title compound was purified by column chromatography (9/1 CHCl₃/MeOH) to give a thick yellow oil. Recrystallization from CH₂Cl₂-Petroleum ether gave a tan solid, mp = 99-101°
- MS (m/e): 410: M⁺, 273 = M⁺-PO₃Et₂
- NMR (CDCl₃) δ = 8.51, 8.50, 7.64 and 7.25 (4m, 4H): aromatic H, 3-picolyl, 6.65 (d, J = 2Hz, 2H): aromatic H, phenyl, 7.75 (broad, 1H): OH, 4.15-3.75 (several m, 5H):


P-O-CH₂-CH₃ and CH-PO₃Et₂, 3.9 (s, 6H): OCH₃, 3.82 and 3.61 (2d, J = 14Hz, 2H): NH-CH₂-Py, 1.31 and 1.16 (2t, J = 7Hz, 6H): P-O-CH₂-CH₃

5 Example 14 - Diethyl α-(3,5-dimethoxy-4-hydroxyphenyl)-N-(2-pyridyl)-amino-methylphosphonate



- A mixture of 3.64 g (20 mmol) of syringaldehyde and 1.88 g (20 mmol) of 2-aminopyridine dissolved in 20 ml toluene and a catalytic amount of TsOH contained in a flask connected to a Dean Stark apparatus was refluxed for 24 h. The solution was evaporated to dryness to give 5.2 g (100%) of the crude imine.
- 10 Diethyl phosphite (5.8 g, 42 mmol) was added to 3.6 g (14 mmol) of the previously described imine dissolved in 25 ml THF and the mixture was refluxed for 20 h. The solvent and the excess of diethyl phosphite were evaporated and the residue was recrystallized from ethanol to give 4.2 g (76%) of a white solid, mp = 163-165°C.
- 15 IR (KBr) = 1240 cm⁻¹ : P=O and 1030 : P-O-C
 MS (m/e) = 397 : M⁺ + 1, 259 (100%) : M⁺ - PO₃Et₂
 NMR (CDCl₃): δ = 8.08, 7.37, 6.60 and 6.41 (4m, 1H each): aromatic H, 2-pyridyl, 6.76 (d, J = 2Hz, 2H): aromatic H, substituted phenyl, 5.6 (s, 1H) : OH, 5.39 (m, 1H): N-H, 5.37 (d x d, 1H, J=9 and 28 Hz): CH-PO₃Et₂, 4.18-3.69 (3m, 4H total): P-O-CH₂-CH₃, 3.87 (s, 6H): OCH₃, 1.24 and 1.15: (2t, J=7Hz): P-O-CH₂-CH₃
- 20


Example 15 - Diethyl α-(3,5-dimethoxy-4-hydroxyphenyl)-N-(4-pyridyl)-aminomethyl phosphonate



- 25 A 25 ml toluene solution containing syringaldehyde (3.64 g, 20 mmol), 4-aminopyridine (1.9 g, 20 mmol) and 5 mg of TsOH placed in a flask connected to a Dean-Stark trap was refluxed for 48 h to give 5.0 g (95%) of the corresponding imine.

- To a suspension of sodium hydride (0.87 g of a 60% mixture, 20 mmol) in 25 ml dry THF was added HPO_3Et_2 (4.14 g, 30 mmol) under nitrogen and the resulting mixture was stirred until the initial turbid suspension became completely clear. To this solution of NaPO_3Et_2 was added the above imine (2.6 g, 10 mmol) dissolved in 5 ml THF and the resulting solution was refluxed for 2 h. THF was evaporated and the residue was partitioned into H_2O and CH_2Cl_2 . Evaporation of the dried organic phase gave a white solid which was recrystallized in EtOH (1.84 g, 45%); mp = 172-174°C.
- MS (m/e) = 396 : M^+ , 259 (100%) : $\text{M}^+ - \text{PO}_3\text{Et}_2$
- NMR (CDCl_3): δ = 8.18, 8.16, 6.48 and 6.46 (4m, 1H each): aromatic H, 4-pyridyl, 6.67 (d, J = 2Hz, 2H): aromatic H, substituted phenyl, 5.27 (d x d, 1H, J = 7 and 10Hz): N-H, 4.66 (d x d, 1H, J = 7 and 23 Hz): $\text{CH}-\text{PO}_3\text{Et}_2$, 4.18-3.60 (3m, 4H total): P-O- CH_2-CH_3 , 3.87 (s, 6H): OCH_3 , 1.30 and 1.15: (2t, J = 7Hz): P-O- CH_2-CH_3


15 Example 16 - Enantiomers of diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(4-picolyl)-aminomethylphosphonate



- a) 3,5-Di-tert-butyl-4-hydroxybenzaldehyde (30 g, 123.5 mmol) and (R)-(+)-1-phenyl-ethylamine (15.7 g, 129.7 mmol) were stirred in 100 ml of THF at room temperature for one day. The solution was dried on MgSO_4 and concentrated. The corresponding imine was recrystallized from ligroin (38 g; 88% yield; mp = 127-128°C).
- The imine (30 g, 89 mmol) and diethylphosphite (15.4 g, 111.3 mmol) were refluxed in 80 ml of toluene for 5 hours. The mixture was evaporated to dryness. HPLC assay of the residue showed that one of the diastereomers is formed predominantly (84% vs 3% of the reaction mixture). The major diastereomer of diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(1-phenyl-ethyl)-aminomethylphosphonate was isolated by successive crystallizations (10g); $[\alpha]_D^{25} + 8.33^\circ$ (c = 1.649, CHCl_3); mp = 105-106°C).
- (+)-Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(1-phenyl-ethyl)-aminomethylphosphonate (9.5 g, 20 mmol) was hydrogenated in ethanol in the presence of 2.5 g of 10% Pd on charcoal to give (-)-diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-aminomethylphosphonate (5.6 g; 76% yield; mp = 143-145°C (recrystallized from ligroin/ CH_2Cl_2); $[\alpha]_D^{25} - 12.12^\circ$ (c = 1.650, CHCl_3).


- (-)-Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-aminomethylphosphonate (11 g, 29.6 mmol) and pyridine-4-carboxaldehyde (6.3 g, 59.3 mmol) were dissolved in 125 ml of MeOH. The mixture was acidified with concentrated HCl (blue bromophenol indicator). After half an hour of stirring at room temperature, NaBH₃CN (5.6 g, 89 mmol) dissolved in 30 ml MeOH was added and the pH was adjusted again with HCl. The reaction mixture was stirred at room temperature for 4 hours then evaporated to dryness and extracted with CH₂Cl₂ and water. The organic phase was dried over MgSO₄ and evaporated. The residue was separated by column chromatography (silicagel, 95/5 CHCl₃/MeOH to give (-)-diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(4-picolyl)-aminomethylphosphonate [(11 g; 80% yield; mp = 66-69°C; $[\alpha]_D^{21}$ -44.05° (c=1.992, CHCl₃)].
- b) 3,5-Di-tert-butyl-4-hydroxybenzaldehyde (30 g, 123.5 mmol) and (S)-(-)-1-phenyl-ethylamine (15.7 g, 129.7 mmol) were stirred in 100 ml of THF for one day to give the corresponding imine (36.5 g; 88% yield; mp = 127-128°C).
- The imine (20 g, 59.3 mmol) and diethylphosphite (10.2 g, 74.2 mmol) were refluxed in 60 ml of toluene for 7 hours. The mixture was evaporated to dryness. HPLC assay of the residue indicated the diastereomeric ratio to be 60 to 40% in addition to starting materials. The latter were stripped off by column chromatography on silicagel (98/2 CH₂Cl₂/MeOH). The fractions containing the mixture of diastereomers were evaporated to dryness and recrystallized three times from ligroin/MTBE to yield the major diastereomer of diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(1-phenyl-ethyl)-aminomethylphosphonate [12 g; mp = 104-105°C; $[\alpha]_D^{28}$ -10.53° (c=1.643, CHCl₃)].
- (-)-Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(1-phenyl-ethyl)-aminomethylphosphonate (42 g, 88.4 mmol) was hydrogenated in ethanol in the presence of 6 g of 10% Pd on charcoal to give (+)-diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-aminomethylphosphonate (24.5 g; 75% yield; mp = 143-144°C (recrystallized from ligroin/MTBE); $[\alpha]_D^{29}$ +11.04° (c=1.714, CHCl₃)).
- (+)-Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-aminomethylphosphonate (11 g, 29.6 mmol) and pyridine-4-carboxaldehyde (6.35 g, 59.3 mmol) in 125 ml of MeOH were reacted with NaBH₃CN (5.6 g, 89 mmol) in the same manner as described for the (-) enantiomer. Column chromatography on silicagel (95/5 CHCl₃ / MeOH) gave (+)-diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(4-picolyl)-aminomethylphosphonate (12 g, 87% yield; mp = 67-70°C; $[\alpha]_D^{21}$ +43.03° (c=1.984, CHCl₃)).

Example 17 - Enantiomers of diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-picolyl)-aminomethylphosphonate



- a) In the same manner as described in example 16, (+)-diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-aminomethylphosphonate (1 g, 2.7 mmol) and pyridine-3-carboxaldehyde (0.43 g, 4 mmol) were reacted with NaBH₃CN (0.34 g, 5.4 mmol) in MeOH for 5 hours at room temperature to yield after trituration in petroleum ether (+)-diethyl- α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-picolyl)-aminomethylphosphonate (1 g, 80% yield; mp = 116-119°C; $[\alpha]_D^{25} +42.88^\circ$ (c=1.614, CHCl₃)).
- b) respectively,
- (-)-diethyl- α -(3,5-di-tert-butyl-4-hydroxyphenyl)-aminomethylphosphonate (1 g, 2.7 mmol) and pyridine-3-carboxaldehyde (0.43 g, 4 mmol) were reacted with NaBH₃CN (0.34 g, 5.4 mmol) in MeOH to give (-)-diethyl- α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-picolyl)-aminomethylphosphonate (0.7 g, 56%; mp = 118-120°C).

Example 18 - Enantiomers of diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate



- The enantiomers of a racemic mixture were separated by preparative HPLC on Chiralcel OD and isocratic elution with hexane/ethanol (9:1), UV detection at 254 nm. Baseline separation was achieved, and the contents of both peaks were evaporated to white solids in which none of the other isomer could be detected by analytical HPLC.

First peak : retention time 18 min, $[\alpha]_D^{20} -7.4^\circ$ (c = 0.244% w/v, EtOH)

Second peak : retention time 34 min, $[\alpha]_D^{20} +8.3^\circ$ (c = 0.255% w/v, EtOH)

The structures of both enantiomers were confirmed by NMR and MS spectroscopies and elemental analysis.

Elemental analysis: $C_{18}H_{25}N_2O_6P$

% Calc. C 54.54 H 6.36 N 7.07

5 (+)-Enantiomer:

mp : 153-157°


% Found C 53.85 H 6.22 N 6.81

(-)-Enantiomer:

mp : 155-158°


10 % Found C 54.25 H 6.24 N 6.94

Example 19 - Enantiomers of diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-phenylpropyl)-aminomethylphosphonate




- 15 a) (-)-Diethyl- α -(3,5-di-tert-butyl-4-hydroxyphenyl)-aminomethylphosphonate (1.7 g, 4.5 mmol) and 3-phenylpropionaldehyde (0.6 g, 4.5 mmol) in 20 ml of absolute methanol were stirred under nitrogen, at room temperature for 30 min. $NaBH_3CN$ (0.3 g, 4.5 mmol) dissolved in 10 ml of methanol was added and the mixture was allowed to react at room temperature for another hour. The reaction mixture was
- 20 evaporated to dryness and the residue dissolved in CH_2Cl_2 . The organic phase was washed with water, then dried over $MgSO_4$. Column chromatography with 98/2 $CHCl_3/MeOH$ as eluent gave (-)-diethyl- α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-phenylpropyl)-aminomethylphosphonate [1.2 g; 56% yield; $[\alpha]_D^{25} +33.1^\circ$ (c=2.055, $CHCl_3$)].
- 25 b) (+) Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-aminomethylphosphonate (1.2 g, 3.2 mmol) and 3-phenylpropionaldehyde (0.4 g, 3.2 mmol) in 20 ml of absolute methanol were reacted in the same manner with $NaBH_3CN$ (0.2 g, 3.2 mmol) in 10 ml of methanol to yield after column chromatography, (+) diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-phenylpropyl)-aminomethylphosphonate [0.94 g; 61%
- 30 yield; $[\alpha]_D^{25} +31.1^\circ$ (c=1.930, $CHCl_3$)].
- c) - The structures of both enantiomers were confirmed by IR, NMR and MS. They were separated by analytical HPLC on Chiralpak AD and isocratic elution with hexane/2-propanol (9:1/v:v).

Example 20 - Diisopropyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-amino-methylphosphonate



- 5 Diisopropyl phosphite (3.3 g, 20 mmol) was added to 2.58 g (10 mmol) of 3,5-dimethoxy-4-hydroxybenzaldehyde N-(3-pyridyl) imine dissolved in 15 ml toluene and the mixture was refluxed for 17 h. The solvent and the excess of diisopropyl phosphite were evaporated and the residue was purified by column chromatography (9/1 CH₂Cl₂/MeOH) and recrystallisation from a mixture of
- 10 EtOH/AcOEt to give 1.56 g (37%) of a white solid, mp = 157-160°.
- MS (m/e) = 424 : M⁺, 259 (100%) : M⁺ - PO₃iPr₂
- NMR (CDCl₃): δ = 8.08, 7.96, 7.03 and 6.84 (4m, 1H each): aromatic H, 3-pyridyl, 6.69 (d, J = 2Hz, 2H): aromatic H, substituted phenyl, 5.8 (broad, 1H) : OH, 4.82 (d x d, 1H, J = 7 and 10Hz): N-H, 4.55 (d x d, 1H, J = 7 and 23Hz): CH-PO₃iPr₂, 4.75-4.65 and 4.55-4.45 (2m, 2H total): P-O-CH-(CH₃)₂, 3.86 (s, 6H): OCH₃, 1.34, 1.28, 1.24 and 0.9: (4d, J = 7Hz): P-O-CH-(CH₃)₂
- 15

Example 21 - Diisopropyl α -(4-hydroxy-3-methoxy-5-methylphenyl)-N-(3-pyridyl)-amino-methylphosphonate




- 20 A mixture of 1.9 g (11 mmol) of 4-hydroxy-3-methoxy-5-methylbenzaldehyde (mp = 98-100°) and 1.08 g (11 mmol) of 3-aminopyridine dissolved in 15 ml toluene and a catalytic amount of p-toluenesulfonic acid (ca. 5 mg) contained in a flask connected to a Dean Stark apparatus was refluxed for 15 h. The solution was evaporated to
- 25 dryness to give 2.7 g (100%) of the crude imine.
- Diisopropyl phosphite (5.48 g, 33 mmol) was added to 2.77 g (11 mmol) of the above imine dissolved in 20 ml THF and the mixture was refluxed for 24 h. The solvent and the excess of diisopropyl phosphite were evaporated and the residue was purified by column chromatography (95/5 CHCl₃/MeOH) and recrystallisation from a

mixture of petroleum ether/CH₂Cl₂ to yield 1.9 g (43%) of a white solid, mp = 123-124°.

MS (m/e) = 408 : M⁺, 243 (100%) : M⁺ - PO₃iPr₂

- 5 NMR (CDCl₃): δ = 8.07, 7.95, 7.02 and 6.84 (4m, 1H each): aromatic H, 3-pyridyl, 6.83- 6.81 : (m, 2H): aromatic H, substituted phenyl, 5.8 (s, 1H) : OH, 4.78 (d x d, 1H, J=7.5 and 10Hz): N-H, 4.30 (d xd, 1H, J=7.5 and 23Hz): CH-PO₃iPr₂, 4.73-4.65 and 4.48-4.40 (2m, 2H total): P-O-CH-(CH₃)₂, 3.85 (s, 3H): OCH₃, 2.22 (s, 3H): CH₃, 1.33, 1.26, 1.24 and 0.96: (4d, J=7Hz): P-O-CH-(CH₃)₂

10 Example 22 - Diisopropyl α-(3-n-butyl-4-hydroxy-5-methoxyphenyl)-N-(3-pyridyl)-amino-methylphosphonate




- A mixture of 6.1 g (30 mmol) of 3-n-butyl-4-hydroxy-5-methoxybenzaldehyde and 2.76 g (30 mmol) of 3-aminopyridine dissolved in 50 ml toluene and a catalytic amount of p-toluenesulfonic acid (ca. 5 mg) contained in a flask connected to a Dean Stark apparatus was refluxed for 16 h. The solution was evaporated to dryness to give 7.8 g (94%) of the crude imine.

- Diisopropyl phosphite (4.20 g, 25 mmol) was added to 2.4 g (8 mmol) of the above imine dissolved in 30 ml THF and the mixture was refluxed for 24 h. The solvent and the excess of diisopropyl phosphite were evaporated and the residue was purified by column chromatography (95/5 CHCl₃/MeOH) and recrystallisation from a mixture of petroleum ether/CH₂Cl₂ to yield 1.9 g (43%) of a white solid, mp = 142-144°.

MS (m/e) = 450 : M⁺, 285 (100%) : M⁺ - PO₃iPr₂

- 25 NMR (CDCl₃): δ = 8.07, 7.95, 7.0 and 6.84 (4m, 1H each): aromatic H, 3-pyridyl, 6.83- 6.80 : (m, 2H): aromatic H, substituted phenyl, 5.8 (s, 1H) : OH, 4.74 (d x d, 1H, J=7.5 and 10Hz): N-H, 4.54 (d xd, 1H, J=7.5 and 23Hz): CH-PO₃iPr₂, 4.75-4.65 and 4.50-4.40 (2m, 2H total): P-O-CH-(CH₃)₂, 3.85 (s, 3H): OCH₃, 2.60 (t, 2H), 1.5 (m, 2H), 1.31 (m, 2H) and 0.90 (t, 3H) : n-Bu, 1.33, 1.26, 1.24 and 0.94: (4d, J=7Hz): P-O-CH-(CH₃)₂
- 30

Example 23 - Diethyl α-(3,5-dimethoxy-4-hydroxyphenyl)-N-methyl-N-(3-picoly)-aminomethylphosphonate




A mixture of 3.0 g (16.5 mmol) of syringaldehyde, 2.03 g (16.6 mmol) of N-methyl-3-picolylamine and 2.3 g (16.6 mmol) diethyl phosphite dissolved in 15 ml toluene and a catalytic amount of p-toluenesulfonic acid (ca. 5 mg) contained in a flask connected to a Dean Stark apparatus was refluxed for 2 h. The solution was evaporated and the residue was purified by column chromatography (95/5 CHCl₃/MeOH) to yield 3.2 g (46%) of a yellow oil.

MS (m/e) = 287 : M⁺ - PO₃Et₂

NMR (CDCl₃): δ = 8.55, 8.51, 7.72 and 7.27 (4m, 1H each): aromatic H, 3-picolyl, 6.73 (d, 2H): aromatic H, substituted phenyl, 5.8 (broad, 1H) : OH, 4.25, 3.94 and 3.7 (3m, 4H) : P-O-CH₂-CH₃, 3.89 (d, J = 23Hz, 1H): CH-PO₃Et₂, 3.9 and 3.4 (2d, 2H) : N(CH₃)-CH₂-Py, 3.91 (s, 6H): OCH₃, 2.41 (s, 3H): N(CH₃)-CH₂-Py, 1.39 and 1.08 (2t, J = 7 Hz, 6 H): P-O-CH₂-CH₃

15 Example 24 - Diisopropyl α-(4-hydroxy-3-methoxy-5-methylphenyl)-N-methyl-N-(3-picolyl)-aminomethylphosphonate



A mixture of 2.0 g (12 mmol) of 4-hydroxy-3-methoxy-5-methylbenzaldehyde, 1.8 g (13.2 mmol) of N-methyl-3-picolylamine and 2.2 g (13.2 mmol) diisopropyl phosphite dissolved in 15 ml toluene and a catalytic amount of p-toluenesulfonic acid (ca. 2 mg) contained in a flask connected to a Dean Stark apparatus was refluxed for 2 h. The solution was evaporated and the residue was purified by column chromatography (95/5 CHCl₃/MeOH) to yield 2.1 g (40%) of a yellow oil.

MS (m/e) = 271 (100%) : M⁺ - PO₃iPr₂

25 NMR (CDCl₃): δ = 8.54, 8.50, 7.72 and 7.24 (4m, 1H each): aromatic H, 3-picolyl, 6.97 and 6.77 (2 m, 2H): aromatic H, substituted phenyl, 5.75 (broad, 1H) : OH, 4.86-4.78 and 4.51-4.42 (2m, 2H total): P-O-CH(CH₃)₂, 3.84 (d, J = 24Hz, 1H): CH-PO₃iPr₂, 3.97 and 3.34 (2d, J = 13.5 Hz, 2H) : N(CH₃)-CH₂-Py, 3.91 (s, 3H):

OCH_3 , 2.36 (s, 3H); CH_3 , 2.26 (s, 3H); $\text{N}(\text{CH}_3)\text{-CH}_2\text{-Py}$, 1.39, 1.37, 1.21 and 0.83 (4d, $J = 7 \text{ Hz}$, 12 H); $\text{P-O-CH}(\text{CH}_3)_2$

The following compounds may also be obtained in an analogous manner to Examples

5 1 to 24:

Diethyl α -(4-hydroxy-3-methoxy-5-n-propylphenyl)-N-(3-pyridyl)-aminomethylphosphonate;


Diisopropyl α -(4-hydroxy-3-methoxy-5-n-propylphenyl)-N-(3-pyridyl)-aminomethylphosphonate;

10 Diethyl α -(3-i-butyl-4-hydroxy-5-methoxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate; and

Diisopropyl α -(3-i-butyl-4-hydroxy-5-methoxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate.

15 Table 1 lists the physicochemical data of compounds of formula (I) that were prepared by the methods illustrated by examples 1-24 of this application. These methods are disclosed in EP 0 559 079A (corresponding to the US Patent 5 424 303).

Table 1 - Aminophosphonates of formula (I)



Cpd	X ¹	X ²	X ³	Z	A	R	mp(°C)	Microanalysis
1	tBu	tBu	H	H	3-pyridyl	Me	168-170	C ₂₂ H ₃₃ N ₂ O ₄ P
2	tBu	tBu	H	H	3-pyridyl	Et	155-156	C ₂₄ H ₃₇ N ₂ O ₄ P
3	tBu	tBu	H	H	3-pyridyl	iPr	135-137	C ₂₆ H ₄₁ N ₂ O ₄ P
4	tBu	tBu	H	H	3-pyridyl	nPr	133-135	C ₂₆ H ₄₁ N ₂ O ₄ P
5	tBu	tBu	H	H	3-pyridyl	nBu	112-114	C ₂₈ H ₄₅ N ₂ O ₄ P
6	tBu	tBu	H	H	4-picolyl	Me	120-122	C ₂₃ H ₃₅ N ₂ O ₄ P
7	tBu	tBu	H	H	4-picolyl	Et	87-91	C ₂₅ H ₃₉ N ₂ O ₄ P
8	tBu	tBu	H	H	4-picolyl	iPr	126-128	C ₂₇ H ₄₃ N ₂ O ₄ P
9	tBu	tBu	H	H	4-picolyl	nPr	108-110	C ₂₇ H ₄₃ N ₂ O ₄ P
10	tBu	tBu	H	H	4-picolyl	nBu	60-61	C ₂₉ H ₄₇ N ₂ O ₄ P
11	tBu	tBu	H	COMe	4-picolyl	Et	160-162	C ₂₇ H ₄₁ N ₂ O ₅ P
12	tBu	tBu	H	H	5-(2-chloropyridyl)	Et	124-126	C ₂₄ H ₃₆ ClN ₂ O ₄ P
13	tBu	tBu	H	H	2-pyridyl	Et	116-118	C ₂₄ H ₃₇ N ₂ O ₄ P
14	tBu	tBu	H	H	4-pyridyl	Et	116-119	C ₂₄ H ₃₇ N ₂ O ₄ P
15	tBu	tBu	H	H	3-picolyl	Et	100-101	C ₂₅ H ₃₉ N ₂ O ₄ P
16	tBu	tBu	H	H	2-picolyl	Et	90-91	C ₂₅ H ₃₉ N ₂ O ₄ P
17	tBu	tBu	H	H	2-(2-pyridyl)ethyl	Et	76-78	C ₂₆ H ₄₁ N ₂ O ₄ P

Table 1 cont

Cpd	X ¹	X ²	X ³	Z	A	R	mp(°C)	Microanalysis
18	H	H	H	H	3-pyridyl	Et	210-212	C ₁₆ H ₂₁ N ₂ O ₄ P
19	OMe	OMe	H	H	3-pyridyl	Me	186-187	C ₁₆ H ₂₁ N ₂ O ₆ P
20	OMe	OMe	H	H	3-pyridyl	Et	181-183	C ₁₈ H ₂₅ N ₂ O ₆ P
21	OMe	OMe	H	H	3-pyridyl	iPr	157-160	C ₂₀ H ₂₉ N ₂ O ₆ P
22	OMe	OMe	H	H	2-picolyl	Et	oil	C ₁₉ H ₂₇ N ₂ O ₆ P
23	OMe	OMe	H	H	3-picolyl	Et	99-101	C ₁₉ H ₂₇ N ₂ O ₆ P
24	OMe	OMe	H	H	4-picolyl	Et	125-127	C ₁₉ H ₂₇ N ₂ O ₆ P
25	OMe	H	H	H	3-pyridyl	Et	171-173	C ₁₇ H ₂₃ N ₂ O ₅ P
26	OEt	H	H	H	3-pyridyl	Et	185-187	C ₁₈ H ₂₅ N ₂ O ₅ P
27	OMe	OMe	Me	H	3-pyridyl	Et	134-136	C ₁₉ H ₂₇ N ₂ O ₆ P
28	Me	Me	H	H	3-pyridyl	Et	176-178	C ₁₈ H ₂₅ N ₂ O ₄ P
29	OMe	OMe	H	H	2-pyridyl	Et	163-165	C ₁₈ H ₂₅ N ₂ O ₆ P
30	OMe	OMe	H	H	4-pyridyl	Et	172-174	C ₁₈ H ₂₅ N ₂ O ₆ P

31	OMe	OMe	H	H	3-pyridyl	nPr	142-143	C ₂₀ H ₂₉ N ₂ O ₆ P
32	OMe	OMe	H	H	3-pyridyl	nBu	158-160	C ₂₂ H ₃₃ N ₂ O ₆ P
33	OMe	NO ₂	H	H	3-pyridyl	Et	212-213	C ₁₇ H ₂₂ N ₃ O ₇ P
34	OMe	OMe	H	H	5-(2-chloropyridyl)	Et	193-195	C ₁₈ H ₂₄ ClN ₂ O ₆ P
35	OMe	OMe	H	H	5-(2-methoxypyridyl)	Et	135-137	C ₁₉ H ₂₇ N ₂ O ₇ P
36	OMe	OMe	H	H	3-(2-methylpyridyl)	iPr	148-150	C ₂₁ H ₃₁ N ₂ O ₆ P
37	OMe	OMe	H	H	5-(2-methylpyridyl)	Et	189-190	C ₁₉ H ₂₇ N ₂ O ₆ P
38	OMe	OMe	H	H	5-(2-methylpyridyl)	iPr	150-152	C ₂₁ H ₃₁ N ₂ O ₆ P
39	Me	t-Bu	H	H	3-pyridyl	Et	90-91	C ₂₁ H ₃₁ N ₂ O ₄ P
40	iPr	iPr	H	H	3-pyridyl	Et	174-176	C ₂₂ H ₃₃ N ₂ O ₄ P
41	sBu	sBu	H	H	3-pyridyl	Et	140-141	C ₂₄ H ₃₇ N ₂ O ₄ P
42	Et	Et	H	H	3-pyridyl	Et	170-171	C ₂₀ H ₂₉ N ₂ O ₄ P
43	OMe	Me	H	H	3-pyridyl	Et	164-166	C ₁₈ H ₂₅ N ₂ O ₅ P
44	OMe	Me	H	H	3-pyridyl	iPr	123-125	C ₂₀ H ₂₉ N ₂ O ₅ P
45	OMe	nBu	H	H	3-pyridyl	Et	133-134	C ₂₁ H ₃₁ N ₂ O ₅ P
46	OMe	nBu	H	H	3-pyridyl	iPr	142-144	C ₂₃ H ₃₅ N ₂ O ₅ P
47	OMe	Me	H	H	3-picolyl	Et	99-101	C ₁₉ H ₂₇ N ₂ O ₅ P
48	OMe	Me	H	H	3-picolyl	iPr	85-86	C ₂₁ H ₃₁ N ₂ O ₅ P
49	OMe	Me	H	H	4-picolyl	Et	129-130	C ₁₉ H ₂₇ N ₂ O ₅ P
50	OMe	Me	H	H	4-picolyl	iPr	138-141	C ₂₁ H ₃₁ N ₂ O ₅ P

Table 1 cont.


Cpd	X ¹	X ²	X ³	Z	A	R	mp(°C)	Microanalysis
51	Me	Me	H	H	3-pyridyl	iPr	169-171	C ₂₀ H ₂₉ N ₂ O ₄ P
52	OEt	H	H	H	3-pyridyl	iPr	192-194	C ₂₀ H ₂₉ N ₂ O ₅ P
53	OEt	Me	H	H	3-pyridyl	Et	172-173	C ₁₉ H ₂₇ N ₂ O ₅ P
54	OEt	Me	H	H	3-pyridyl	iPr	177-178	C ₂₁ H ₃₁ N ₂ O ₅ P
55	OEt	OEt	H	H	3-pyridyl	Et	130-132	C ₂₀ H ₂₉ N ₂ O ₆ P
56	OEt	OEt	H	H	3-pyridyl	iPr	149-150	C ₂₂ H ₃₃ N ₂ O ₆ P
57	OMe	Et	H	H	3-pyridyl	Et	139-141	C ₁₉ H ₂₇ N ₂ O ₅ P
58	OMe	Et	H	H	3-pyridyl	iPr	146-148	C ₂₁ H ₃₁ N ₂ O ₅ P
59	OMe	OEt	H	H	3-pyridyl	Et	156-157	C ₁₉ H ₂₇ N ₂ O ₆ P
60	OMe	OEt	H	H	3-pyridyl	iPr	159-160	C ₂₁ H ₃₁ N ₂ O ₆ P
61	OMe	OMe	H	Me	3-picoyl	Et	oil	*NMR and MS
62	OMe	OMe	H	Me	3-picoyl	iPr	oil	*NMR and MS
63	OMe	Me	H	Me	3-picoyl	Et	oil	*NMR and MS
64	OMe	Me	H	Me	3-picoyl	iPr	oil	*NMR and MS
65	OMe	OMe	H	H	3-pyridyl	Et	153-157	**C ₁₈ H ₂₅ N ₂ O ₆ P
66	OMe	OMe	H	H	3-pyridyl	Et	155-158	***C ₁₈ H ₂₅ N ₂ O ₆ P
67	OMe	OMe	H	H	phenyl	Et	143-146	C ₁₉ H ₂₆ NO ₆ P
68	OMe	OMe	H	H	phenyl	iPr	144-146	C ₂₁ H ₃₀ NO ₆ P
69	OMe	Me	H	H	5-(2-methylpyridyl)	Et	154-155	C ₁₉ H ₂₇ N ₂ O ₅ P
70	OMe	Me	H	H	5-(2-methylpyridyl)	iPr	149-150	C ₂₁ H ₃₁ N ₂ O ₅ P
71	OMe	Me	H	H	3-(2-methylpyridyl)	Et	150-152	C ₁₉ H ₂₇ N ₂ O ₅ P
72	OMe	Me	H	H	3-(2-methylpyridyl)	iPr	148-150	C ₂₁ H ₃₁ N ₂ O ₅ P
73	OMe	OMe	H	H	3-(2-methylpyridyl)	Et	146-148	C ₁₉ H ₂₇ N ₂ O ₆ P

* : Identified by NMR and MS spectroscopies

5 ** : (+)Enantiomer of Compound 20

*** : (-)Enantiomer of Compound 20

Table 1 - Aminophosphonates of formula (I), (cont.)



Cpd	X ¹	X ²	X ³	(B) _n	Z	R	mp(°C)	Microanalysis
	H	O	CH ₂	bond	H	Et	98-99	C ₁₇ H ₂₁ N ₂ O ₅
	OMe	OMe	H	CH=CH	H	Et	foam	*NMR and MS
	OMe	OMe	H	CH ₂ -CH ₂	H	Et	134-138	C ₂₀ H ₂₉ N ₂ O ₆ P

5 * : Identified by NMR and MS spectroscopies

Biological Data

10 In vitro Data - The compounds of formula (I) were tested for lowering the production of Lp(a) in primary cultures of Cynomolgus hepatocytes according to the assays described below. Two incubation times were used :4 h for Assay 1 and 24 h for Assay 2.

15 Protocol - Hepatocytes were isolated from livers of adult Cynomolgus monkeys by the two-step collagenase perfusion method according to C. Guguen-Guillouzo and A. Guillouzo "Methods for preparation of adult and fetal hepatocytes" p.1-12 in "Isolated and Cultured Hepatocytes", les editions Inserm Paris and John Libbey Eurotext London (1986).

20 The viability of cells was determined by Trypan blue staining. The cells were then seeded at a density of 1.5-2.10⁵ viable cells per 2cm² in 24 well tissue culture plates in a volume of 500µl per well of Williams E tissue culture medium containing 10% fetal calf serum. Cells were incubated for 4-6 hours at 37°C in a CO₂ incubator (5% CO₂) in the presence of 20µM of the test compounds dissolved in ethanol. Four wells were used for each compound. Nicotinic acid and steroid hormones were used as references to validate the assay system since they are known to decrease Lp(a) in man. Control cells were incubated in the presence of ethanol only.

25 The amount of Lp(a) secreted in culture medium was directly assayed by ELISA using a commercially available kit. Cells were washed and lysed as described by A.L. White et al, Journal of Lipid Research vol 34, p. 509-517, (1993) and the cellular content of Lp(a) was assayed as described above.

Changes in Lp(a) concentration in culture medium are given as the percentage of value measured for the control plates at 4 h (Assay 1) or 24 h (Assay 2).

Results - Assay 1: compounds 2, 7, 11, 15, 16, 18 and 20 were found to change the concentrations of Lp(a) in the culture medium in the range from -12 to -34%.

- 5 **Assay 2:** compounds 1, 2, 3, 5, 7, 11, 13, 15, 17, 19, 20, 21, 26 to 29, 32, 34 to 52, 57 to 60, 65 and 66 were found to change the concentrations of Lp(a) in the culture medium in the range from -7 to -37%.

- 10 ***In Vivo* Data - Study Protocol** - Male cynomolgus monkeys weighing between 3 and 7 kg were divided into groups of 3 to 4 animals each. Prior to treatment their plasma Lp(a) levels were followed over a two month period to ascertain a constant baseline value. Test compounds were given orally by gavage at the dose of 25 mg/kg/day for 4 weeks and Lp(a) was measured at day 28. At the end of the dosing period, animals were maintained for a treatment free period of 4 weeks, whereupon their plasma
- 15 Lp(a) levels returned to pretreatment levels. This control provided proof that the decrease in Lp(a) measured was caused by the pharmacological activity of the test compounds.

Results - At Days -7 and 28, after an overnight fast, blood samples were collected on EDTA and Lp(a) was measured by the highly sensitive and specific ELISA test.


- 20 Results (mean of 3-4 values of each group) were expressed as % of predose (Day -7). Selected compounds of formula (I) were tested under the experimental conditions to investigate their pharmacological activity *in vivo*.

- 25 The compounds No 1, 2, 3, 7, 15, 17, 19, 20, 21, 27, 28, 32, 39, 44 and 52 lower plasma Lp(a) in the range of -13% to -51% (value measured at Day 28, % change from predose at Day -7).

- 30 The compounds of Formula (I) have therefore a therapeutic potential for the treatment of the following diseases where Lp(a) is associated with accelerated atherosclerosis, abnormal proliferation smooth muscle cells and increased thrombogenesis :coronary heart disease, peripheral artery disease : intermittent claudication, extracranial carotid atherosclerosis, stroke, restenosis after angioplasty and atherosclerosis occurring after heart transplant. The primary indications of these compounds would be the treatment of the diseases mentioned above.

Claims

1. The use of a compound of formula (I):



5 where:

X^1 , X^2 , identical or different, are H, a straight or branched alkyl or alkoxy group having from 1 to 8 carbon atoms, a hydroxy group or a nitro group,

X^3 is H, an alkyl group from 1 to 4 carbon atoms, X^3O and one of the two other substituents X^1 or X^2 may form an alkylidene dioxy ring having from 1 to 4 carbon atoms,

R^1 , R^2 , identical or different, are H, a straight or branched alkyl group having from 1 to 6 carbon atoms,


B is CH_2 , CH_2-CH_2 or $CH=CH$,

n is zero or 1,

15 Z is H, a straight or branched alkyl group having from 1 to 8 carbon atoms, an acyl group R^3-CO where R^3 is an alkyl group from 1 to 4 carbon atoms, a perfluoroalkyl group from 1 to 4 carbon atoms,

A is H, $CH_2-CH=CH_2$, a straight, branched or cyclic alkyl group having from 1 to 8 carbon atoms, or is selected from the following groups :

20




where :

- 5 k is an integer from 2 to 4, m is 0 or an integer from 1 to 5, X^4 , X^5 , X^6 , identical or different, are H, a straight or branched alkyl or alkoxy group from 1 to 8 carbon atoms, a hydroxy, trifluoromethyl, nitro, amino, dimethylamino, diethylamino group, a halogen atom (F, Cl, Br, I), X^4 and X^5 may form an alkylidendioxy ring having from 1 to 4 carbon atoms, X^7 is H or CH_3 , R is a straight or branched alkyl group

- having from 1 to 6 carbon atoms, an aryl or arylalkyl group from 6 to 9 carbon atoms,
 or a pharmaceutically acceptable salt thereof;
 in the manufacture of a medicament for use in decreasing plasma and tissue
 5 lipoprotein(a) levels.
2. A use according to claim 1 for the manufacture of a medicament for the treatment
 of thrombosis by decreasing plasma lipoprotein(a) levels.
- 10 3. A use according to claim 1 for the manufacture of a medicament for the treatment
 of restenosis following angioplasty by decreasing plasma lipoprotein(a) levels.
4. A use according to claim 1 for the manufacture of a medicament for the treatment
 of atherosclerosis by decreasing plasma lipoprotein(a) levels.
- 15 5. A use according to any one of claims 1 to 4 wherein the compound of formula (I)
 is selected from:
 diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(4-picolyl)-
 aminomethylphosphonate,
 20 diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(2-picolyl)-
 aminomethylphosphonate,
 diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-picolyl)-
 aminomethylphosphonate,
 diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-[2-(2-pyridyl)ethyl]-
 25 aminomethylphosphonate,
 diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-pyridyl)-
 aminomethylphosphonate,
 dimethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-pyridyl)-
 aminomethylphosphonate,
 30 diisopropyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-pyridyl)-
 aminomethylphosphonate,
 diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(2-pyridyl)-
 aminomethylphosphonate,
 diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(4-pyridyl)-
 35 aminomethylphosphonate,
 diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-[5-(2-chloropyridyl)]-
 aminomethylphosphonate.

- diethyl α -(4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
 diethyl α -(3,4-methylenedioxyphenyl)-N-(3-pyridyl)-aminophosphonate,
 diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
 dimethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-
 5 aminomethylphosphonate,
 diisopropyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-
 aminomethylphosphonate,
 diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(2-pyridyl)-aminomethylphosphonate,
 diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(4-pyridyl)-aminomethylphosphonate,
 10 diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(2-picolyl)-aminomethylphosphonate,
 diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-picolyl)-aminomethylphosphonate,
 diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(4-picolyl)-aminomethylphosphonate,
 diethyl α -(4-hydroxy-3-methoxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
 diethyl α -(3-ethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
 15 diethyl α -(3,4,5-trimethoxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
 diethyl α -(3,5-dimethyl-4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
 (+)-diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(4-picolyl)-
 aminomethylphosphonate,
 (-)-diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(4-picolyl)-
 20 aminomethylphosphonate,
 (+)-diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-picolyl)-
 aminomethylphosphonate,
 (-)-diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-picolyl)-
 aminomethylphosphonate,
 25 (+)-diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-
 aminomethylphosphonate,
 (-)-diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-
 aminomethylphosphonate,
 (+)-diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-phenylpropyl)-
 30 aminomethylphosphonate,
 (-)-diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-phenylpropyl)-
 aminomethylphosphonate,
 diisopropyl α -(3,5-di-methoxy-4-hydroxyphenyl)-N-[5-(2-methylpyridyl)]-
 aminomethylphosphonate,
 35 diethyl α -(3-tert-butyl-4-hydroxy-5-methylphenyl)-N-(3-pyridyl)-amino-
 methylphosphonate,

- diethyl α -(4-hydroxy-3-methoxy-5-methylphenyl)-N-(3-pyridyl)-amino-
methylphosphonate,
diisopropyl α -(4-hydroxy-3-methoxy-5-methylphenyl)-N-(3-pyridyl)-amino-
methylphosphonate,
5 diethyl α -(3-n-butyl-4-hydroxy-5-methoxyphenyl)-N-(3-pyridyl)-amino-
methylphosphonate,
diisopropyl α -(3-n-butyl-4-hydroxy-5-methoxyphenyl)-N-(3-pyridyl)-amino-
methylphosphonate,
diethyl α -(4-hydroxy-3-methoxy-5-n-propylphenyl)-N-(3-pyridyl)-
10 aminomethylphosphonate;
diisopropyl α -(4-hydroxy-3-methoxy-5-n-propylphenyl)-N-(3-pyridyl)-
aminomethylphosphonate;
diethyl α -(3-i-butyl-4-hydroxy-5-methoxyphenyl)-N-(3-pyridyl)-
aminomethylphosphonate;
15 diisopropyl α -(3-i-butyl-4-hydroxy-5-methoxyphenyl)-N-(3-pyridyl)-
aminomethylphosphonate,
Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-pyridyl)
aminomethylphosphonate;
Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(2-picolyl)
20 aminomethylphosphonate;
Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-picolyl)
aminomethylphosphonate;
Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-methyl-N-(3-picolyl)
aminomethylphosphonate;
25 Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(2-pyridylethyl)
aminomethylphosphonate; and
Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(2-picolyl)
aminomethylphosphonate.
- 30 6. A use as claimed in any one of claims 1 to 4 in which the compound of formula
(I) is represented by the sub-formula (Ia):



or a pharmaceutically acceptable salt thereof.

7. A use as claimed in claim 6 in which:

X^1 is H, $C_{(1-8)}$ alkyl or $C_{(1-8)}$ alkoxy;

5 X^2 is $C_{(1-8)}$ alkyl or $C_{(1-8)}$ alkoxy;

X^3 is H, $C_{(1-4)}$ alkyl, or X^3O and one of the two other substituents X^1 or X^2 may form an alkylidene dioxy ring having from 1 to 4 carbon atoms;

R^1 , R^2 , which may be identical or different, are H or $C_{(1-6)}$ alkyl;

B is CH_2CH_2 , $CH=CH$, or CH_2 ;

10 n is zero or 1;

Z is H or a $C_{(1-8)}$ alkyl group;

m is 0 or an integer from 1 to 5;

X^4 is H, or $C_{(1-8)}$ alkyl, $C_{(1-8)}$ alkoxy or halo;

and the pyridyl ring is attached by the ring carbon α - or β - to the nitrogen (2- or 3-

15 pyridyl);

or a salt, preferably a pharmaceutically acceptable salt, thereof.

8. A use as claimed in claim 7 in which, in the compound of formula (Ia), X^1 is H, $C_{(1-4)}$ alkyl or $C_{(1-4)}$ alkoxy.

20

9. A use as claimed in claim 8 in which, in the compound of formula (Ia), X^1 is hydrogen, methyl or methoxy.

10. A use as claimed in any one of claims 7 to 9 in which, in the compound of formula (Ia), X^2 is $C_{(1-4)}$ alkyl or $C_{(1-4)}$ alkoxy.

25

11. A use as claimed claim 10 in which, in the compound of formula (Ia), X^2 is methyl or methoxy.

12. A use as claimed in claim 7 in which, in the compound of formula (Ia), X^1 and X^2 are both alkoxy or one of X^1 and X^2 is alkyl and the other is alkoxy, or one of X^1 and X^2 is $C_{(1-4)}$ alkyl and the other of X^1 and X^2 is $C_{(1-3)}$ alkyl.

30

13. A use as claimed in claim 12 in which, in the compound of formula (Ia), X^1 and X^2 are methoxy and methoxy, methoxy and methyl, n-propyl or iso-butyl, or methyl and methyl or t-butyl, respectively.

35

14. A use as claimed in any one of claims 7 to 13 in which, in the compound of formula (Ia), X^3 is hydrogen.

40


15. A use as claimed in any one of claims 7 to 14 in which, in the compound of formula (Ia), $(B)_n$ is a direct bond.

16. A use as claimed in any one of claims 7 to 15 in which, in the compound of formula (Ia), R^1 and R^2 is each a straight or branched $C_{(1-3)}$ alkyl group.
- 5 17. A use as claimed in claim 16 in which, in the compound of formula (Ia), R^1 and R^2 is each a C_2 or C_3 alkyl group.
18. A use as claimed in any one of claims 7 to 17 in which, in the compound of formula (Ia), Z is hydrogen.
- 10 19. A use as claimed in any one of claims 7 to 18 in which, in the compound of formula (Ia), X^4 is hydrogen or methyl which is preferably on the ring carbon adjacent to N.
- 15 20. A use as claimed in any one of claims 7 to 19 in which, in the compound of formula (Ia), the pyridyl ring is attached by the ring carbon β - to the nitrogen (3-pyridyl).
21. A compound of formula (Ia) as defined in any one of claims 7 to 20 and
 20 excluding:
 Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-pyridyl)
 aminomethylphosphonate;
 Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(2-picolyl)
 aminomethylphosphonate;
 25 Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-picolyl)
 aminomethylphosphonate;
 Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-methyl-N-(3-picolyl)
 aminomethylphosphonate;
 Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(2-pyridylethyl)
 30 aminomethylphosphonate; and
 Diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(2-picolyl)
 aminomethylphosphonate.
22. A compound of formula (Ia) as defined in claim 21 selected from:
 35 dimethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-pyridyl)-
 aminomethylphosphonate,
 diisopropyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-pyridyl)-
 aminomethylphosphonate,
 diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(2-pyridyl)-
 40 aminomethylphosphonate,

- diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(4-pyridyl)-aminomethylphosphonate,
 diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-[5-(2-chloropyridyl)]-aminomethylphosphonate.
- 5 diethyl α -(4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
 diethyl α -(3,4-methylenedioxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
 diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
 dimethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
- 10 diisopropyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
 diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(2-pyridyl)-aminomethylphosphonate,
 diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(4-pyridyl)-aminomethylphosphonate,
 diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(2-picolyl)-aminomethylphosphonate,
- 15 diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-picolyl)-aminomethylphosphonate,
 diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(4-picolyl)-aminomethylphosphonate,
 diethyl α -(4-hydroxy-3-methoxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
 diethyl α -(3-ethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
 diethyl α -(3,4,5-trimethoxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
- 20 diethyl α -(3,5-dimethyl-4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
 (+)-diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(4-picolyl)-aminomethylphosphonate,
 (-)-diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(4-picolyl)-aminomethylphosphonate,
- 25 (+)-diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-picolyl)-aminomethylphosphonate,
 (-)-diethyl α -(3,5-di-tert-butyl-4-hydroxyphenyl)-N-(3-picolyl)-aminomethylphosphonate,
 (+)-diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
- 30 (-)-diethyl α -(3,5-dimethoxy-4-hydroxyphenyl)-N-(3-pyridyl)-aminomethylphosphonate,
 diisopropyl α -(3,5-di-methoxy-4-hydroxyphenyl)-N-[5-(2methyl-pyridyl)]-aminomethylphosphonate,
- 35 diethyl α -(3-tert-butyl-4-hydroxy-5-methylphenyl)-N-(3-pyridyl)-aminomethylphosphonate,

- diethyl α -(4-hydroxy-3-methoxy-5-methylphenyl)-N-(3-pyridyl)-amino-
methylphosphonate,
diisopropyl α -(4-hydroxy-3-methoxy-5-methylphenyl)-N-(3-pyridyl)-amino-
methylphosphonate,
5 diethyl α -(3-n-butyl-4-hydroxy-5-methoxyphenyl)-N-(3-pyridyl)-amino-
methylphosphonate,
diisopropyl α -(3-n-butyl-4-hydroxy-5-methoxyphenyl)-N-(3-pyridyl)-amino-
methylphosphonate,
diethyl α -(4-hydroxy-3-methoxy-5-n-propylphenyl)-N-(3-pyridyl)-
10 aminomethylphosphonate;
diisopropyl α -(4-hydroxy-3-methoxy-5-n-propylphenyl)-N-(3-pyridyl)-
aminomethylphosphonate;
diethyl α -(3-i-butyl-4-hydroxy-5-methoxyphenyl)-N-(3-pyridyl)-
aminomethylphosphonate; and
15 diisopropyl α -(3-i-butyl-4-hydroxy-5-methoxyphenyl)-N-(3-pyridyl)-
aminomethylphosphonate.

23. A pharmaceutical composition comprising a compound of formula (Ia) as defined
in claim 21 and a pharmaceutically acceptable carrier or excipient.
20
24. A compound of formula (Ia) as defined in claim 21 for use in therapy.
25. A process for preparing a compound of formula (Ia) as defined in claim 21 which
process comprises
25 (a) when Z is hydrogen, treating an imine of formula (II):



(II)

- in which B, X¹, X², X³ and n are as defined in claim 21;
30 with a phosphite compound of formula (III):




(III)

in which R^1 and R^2 are as defined in claim 21; or a trialkyl silyl derivative or metal salt thereof;

in the presence or absence of a catalyst, optionally in a solvent;

- 5 (b) for compounds of formula (Ia) in which Z is not hydrogen, treating equimolar amounts of an aldehyde of formula (V):



(V)

- 10 in which B, X^1 , X^2 , X^3 and n are as defined in claim 21; with a secondary amine of formula (VII):




(VII)

- 15 in which Z is a $C_{(1-8)}$ alkyl group and A is as hereinbefore defined; and a phosphite of formula (III):


(c) in compounds of formula (I) in which m is not zero, treating a compound of formula (VIII):

20



(VIII)

in which B, R^1 , R^2 , X^1 , X^2 , X^3 and n are as defined in claim 21; an aldehyde of formula (IX):




25

(IX)

in which m is an integer from 1 to 5 and X^4 is as hereinbefore defined under reductive amination conditions.

26. A process for preparing an individual enantiomer of an aminophosphonate of formula (I) which process comprises treating either of the (+) or (-) enantiomer of the α -substituted aminomethylphosphonate of formula (X):



(X)

in which B, R^1 , R^2 , X^1 , X^2 , X^3 and n are as defined in claim 1;
with an aldehyde of formula (XI):



(XI)

in which R^3 is as defined in claim 1;
under reductive amination conditions.

27. A process as claimed in claim 26 in which the reaction is carried out in the presence of sodium cyanoborohydride in an alcoholic solvent, preferably methanol, at a pH between 3 to 6 and at a temperature between $0^\circ C$ and $25^\circ C$.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 96/02842

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/66 A61K31/675

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 559 079 (SYMPHAR S.A.) 8 September 1993 cited in the application see the whole document & ZH.OBSHCH. KHIM., vol. 51, no. 2, 1981, pages 341-348, A.S.REMIZOV ET AL.: ---	1,21-27
X,P	EP,A,0 703 239 (HOECHST AKTIENGESELLSCHAFT) 27 March 1996 see the whole document --- -/--	21-27

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

23 September 1996

Date of mailing of the international search report

15. 10. 96

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Theuns, H

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 96/02842

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>CHEMICAL ABSTRACTS, vol. 95, no. 17, 26 October 1981 Columbus, Ohio, US; abstract no. 150763b, XP002013995 see abstract & ZH.OSHCH.KHIM., vol. 51, no. 2, 1981, pages 341-348, -----</p>	21-27

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 96/02842

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 1,5-20
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
The expression "decreasing plasma and tissue lipoprotein (a) levels" is not a proper description of a therapeutic application, because it is not immediately clear which disorders may be treated as a result of such.

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 96/02842

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0559079	08-09-93	CH-A- 683996	30-06-94
		AU-A- 3394793	09-09-93
		CA-A- 2091031	06-09-93
		JP-A- 6049083	22-02-94
		NZ-A- 247056	28-08-95
		US-A- 5424303	13-06-95
		ZA-A- 9301473	23-09-93

EP-A-0703239	27-03-96	DE-A- 4433244	28-03-96
		AU-A- 3068695	04-04-96
		CA-A- 2158517	20-03-96
		JP-A- 8104694	23-04-96
